Autism Diagnostic Observation Schedule

Purpose: To Evaluate individuals suspected of having characteristics

similar to Autism and other diagnoses along the Pervasive Developmental

(PDD) spectrum.

➤ Age Range: Toddlers to Adults

> Contains: 4 Modules

Time to Administer: Each module requiring just 35 to 40 minutes

Uses Standardized materials.

Autism Screening Instrument for Educational Planning Second Edition (ASIEP-2)

Published: 1978 to 1993

> Purpose: A comprehensive list of screening tools and sub-tests which all screen or evaluate

Autistic features in children and adults.

> Scores: Sensory, Relating Body and Object Use; Language, Social and Self Help; Autism

Behavior, vocal Behavior, Repetitive, Non-communicative, Babbling, Unintelligible, Interaction, Independent Play, No Response, Negative, Educational Assessment, In Seat, Receptive Language, Expressive Language, Body Concept, Speech Imitation, Hand-Shaping, Random Position-A, Fixed Position-B, Fixed Position-C, Random

Position-D.

➤ 5 Sub-tests: Autism Behavior Checklist, The Sample of Vocal Behavior, The Interaction assessment, the educational assessment, and the prognosis of learning rate.

➤ Age Range: 18 months to 49 years.

> Uses standardized materials and age referenced materials. Also utilizes curriculum-based and criterion referenced materials.

Autism Screening Questionnaire/Social Communication Questionnaire (SCQ)

➤ Published: Formally ASQ, recently reformatted in 1999

> Purpose: To screen for Autism and

➤ Age Range: 4 to adult

> Contains:

> Time to Administer:

> Zero to Three Caution: The SCQ has primarily been tested with older children (six years and above)

and with parents who had already received a full ADI-R. Therefore, results with a younger population or with parents less familiar with these kinds of

questions may be less clear.

> Uses standardized and age referenced materials.

Behavior Rating Instrument for Autistic and other atypical children (BRIAC)

> Purpose: Assesses seven scales of development for Autistic children

> Age Range: Autistic children of all ages.

Seven scales measuring the following areas: Relationship to an adult; > Contains:

> Communication; Drive for Mastery; Vocalization and Expressive Speech; Sound and Sound Reception; Social Responsiveness; and Psychobiological Development. Two additional scales assess non-vocal communication for children who use

manual communication due to vocal and/or hearing deficits.

> Uses Standardized materials.

Checklist for Autism in Toddlers (CHAT)

Published: 1992

Purpose: To Screen toddlers at risk for social and communication disorders

Age Range: 18 months and above

Contains: Parent Report and Observation items. Toddlers are scored using pretend

play, following a point; proto-declarative pointing; pretending and producing a point.

Time to Administer: 5-10 min.

Uses Standardized pass or fail ratings.

Childhood Autism Rating Scale Standardized (CARS)

Published:

This Scale helps to identify children with Autism and to distinguish them from Purpose:

developmentally disabled children without diagnoses of Autism. It also

distinguishes mid-to-moderate from severe autism.

Age Range: Suitable for use with any child over 2 years of age.

Contains: This Scale includes a 15 item behavior-rating scale.

Time to Administer: 5 to 10 minutes.

Uses Criterion or cut scores as its norms.

Computerized FAS Facial Photographic Screening Tool

Identifies 3 facial features of FAS: small eyes, smooth philtrum and a thin upper > Purpose:

➤ Sensitivity100%

> Specificity: 99.8%

➤ Predictive Value Positive: 83.3% ➤ Predictive Value Negative: 100%

Devereux Early Childhood Assessment Program (DECA)

Published: 1998

Purpose: To identify children who may be at risk for low attachment, self control, initiative,

and may struggle with behavior. Also assesses classroom strategies as well as

effective professional practice with families.

Age Range: 24 months and above, however, birth to three edition is currently pending.

> Contains: Contains four separate Scales

> Time to Administer:

> Spanish version available.

Uses Standardized ratings and age referenced materials.

Early and Periodic Screening, Diagnostic, and Treatment (EPSDT)

Statewide implementation: 1995

- Two mutually supportive, operation components: assuring the availability and accessibility and accessibility of required health care resources; and helping Medicaid recipients and their parents or guardians effectively use these resources.
- ❖ Must include the following services: screening services: comprehensive health and developmental history, comprehensive unclothed physical exam, appropriate immunizations, laboratory tests, health education, vision services, dental services, hearing services, and other necessary health care.
 - Diagnosis
 - Treatment
 - Lead poisoning prevention
- **State Medicaid Agency Required activities:**
 - Inform all Medicaid eligible persons under 21 that EPSDT services are available;
 - set distinct periodicity schedules for screening, dental, vision, and hearing services;
 - report EPSDT performance information annually.
- **EPSDT** mandated treatment services:
 - Vision care
 - Hearing care
 - Dental care
 - Podiatrist services
 - Optometrist services
 - Chiropractor services
 - Physician services
 - Medical and remedial care
 - Home health services
 - Private duty nursing services
 - Clinic services furnished under physician direction
 - Nursing facility services
 - Inpatient hospital care
 - Outpatient hospital care
 - Personal care services

Early and Periodic Screening, Diagnostic, and Treatment (EPSDT) Continued

- Transportation
- Case management
- Hospice care
- Preventive services
- Federally qualified health center and rural health clinic services
- Family planning services
- Laboratory and x-ray services
- Emergency hospital services
- Rehabilitation services
- Intermediate care facilities
- Intermediate care facilities services for the mentally retarded
- Impatient psychiatric services
 - Christian science nurses/sanatoria
 - Physical therapy and related services
 - Occupational therapy
 - Speech and language care
 - Prescribed drugs, dentures, and prostheses
 - Nurse midwife services
 - Respiratory care
 - Certified pediatric and family nurse practitioner services
 - Community supported living arrangements for persons with developmental disabilities
 - Other diagnostic, screening, preventative, and medical or remedial services

The FAS Screen: A Rapid Screening Tool for Fetal Alcohol Syndrome

- ➤ *Purpose*: Designed for use in a community setting and in the clinical setting by both paraprofessional and medically trained personnel.
- > Age Range: 4 to 18 Years.
- Contains: Includes screening items for growth impairment, neuralgic dysfunction and facial features, mental retardation, common behavioral signs present in diagnosed cases of FAS.
- ➤ *Needed Training:* 4 hours
- ➤ Time to Administer: 10 to 15 Minutes.
- > Specificity: 94.1%: did not have sufficient signs to be diagnosed with FAS.
- ➤ Positive predictive value: 9.1%

Gilliam Autism Rating Scale (GARS)

➤ Published: 1995

Age Range: 3 to 22 years of age.

Administration: Administered by teacher, teacher's aide, or parent to a group of children.

➤ Time to Administer: 5 to 10 minutes

> Consists: Behavioral Checklist with 4 sub-scales consisting of Stereotyped

Gilliam Autism Rating Scale (GARS) CONTINUED

Behaviors, Communication, Social Interaction, & Developmental

Disturbances

> Scores: Stereotyped Behaviors, Communication, Social Interaction,

Developmental, and Autism Quotient

➤ Uses Standardized scores and Norm-referenced materials.

Modified Checklist for Autism in Toddlers (M-CHAT)

> Age Range: 18 to 24 Months

➤ Purpose: Modified from the original CHAT from the UK for sensitivity for American

children.

➤ Contains: 23 yes/no questions aimed at screening for autistic spectrum disorders.

Uses Standardized age referenced materials.

Parent Interviews for Autism (PIA)

➤ A set of structured questions regarding the child's developmental progress. PIA's are often used in conjunction with other observational tools to evaluate the child fully.